European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC)

Fourth Report by the United Kingdom under Article 17

on the implementation of the Directive from January 2013 to December 2018

Supporting documentation for the conservation status assessment for the species:

S1095 - Sea lamprey (Petromyzon marinus)

SCOTLAND

IMPORTANT NOTE - PLEASE READ

- The information in this document is a country-level contribution to the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- The 2019 Article 17 UK Approach document provides details on how this supporting information was used to produce the UK Report.
- The UK Report on the conservation status of this species is provided in a separate document.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Explanatory notes (where provided) by the country are included at the end. These provide an audit trail of relevant supporting information.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species) and/or (iv) the field was only relevant at UK-level (sections 9 Future prospects and 10 Conclusions).
- For technical reasons, the country-level future trends for Range, Population and Habitat for the species are only available in a separate spreadsheet that contains all the country-level supporting information.
- The country-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, https://jncc.gov.uk/article17, for further information on UK Article 17 reporting.

| NATIONAL LEVEL | | |
|-----------------------------------------|--------------------------------|--|
| 1. General information | | |
| 1.1 Member State | UK (Scotland information only) | |
| 1.2 Species code | 1095 | |
| 1.3 Species scientific name | Petromyzon marinus | |
| 1.4 Alternative species scientific name | | |
| 1.5 Common name (in national language) | Sea lamprey | |

2. Maps

| 2.2 Year or period 1990-2018 |
|----------------------------------------------------------------------------------------------|
| |
| 2.3 Distribution map Yes |
| 2.4 Distribution map Method used Based mainly on extrapolation from a limited amount of data |
| 2.5 Additional maps No |

3. Information related to Annex V Species (Art. 14)

| 3. Illiorillation related to | Allilex v Species (Alt. 14) | |
|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|----|
| 3.1 Is the species taken in the wild/exploited? | No | |
| 3.2 Which of the measures in Art. 14 have been taken? | a) regulations regarding access to property | No |
| | b) temporary or local prohibition of the taking of specimens in the wild and exploitation | No |
| | c) regulation of the periods and/or methods of taking specimens | No |
| | d) application of hunting and fishing rules which take account of the conservation of such populations | No |
| | e) establishment of a system of licences for taking specimens or of quotas | No |
| | f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens | No |
| | g) breeding in captivity of animal species as well as artificial propagation of plant species | No |

h) other measures

No

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

| b) Statistics/ quantity taken | Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period | | | | | |
|----------------------------------|-----------------------------------------------------------------------------------------------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Season/ year 1 | Season/ year 2 | Season/ year 3 | Season/ year 4 | Season/ year 5 | Season/ year 6 |
| Min. (raw, ie. not rounded) | | | | | | |
| Max. (raw, ie. not rounded) | | | | | | |
| Unknown | No | No | No | No | No | No |

- 3.4. Hunting bag or quantity taken in the wild Method used
- 3.5. Additional information

BIOGEOGRAPHICAL LEVEL

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

4.2 Sources of information

Atlantic (ATL)

Maitland PS. 2000. Distribution of lampreys in the River Teith. Unpublished Scottish Natural Heritage commissioned report.

APEM. 2004. Distribution of sea, brook and river lampreys on the River Tay. Scottish Natural Heritage Commissioned Report No. 032.

Ecological Research Associates. 2017. National lamprey survey of Scotland. Scottish Natural Heritage Commissioned Report No. 464.

The Tweed Foundation. Assessment of lamprey distribution and abundance in the River Tweed cSAC/SSSI 2004. Unpublished Scottish Natural Heritage commissioned report.

Watt J, Ravenscroft NOM, Seed M. 2008. Site Condition Monitoring of lamprey in the River Tay SAC. Scottish Natural Heritage Commissioned Report No. 292. Watt J, Brown L, Bull, C. Lamprey Site Condition Monitoring of the River Spey SSSI and SAC 2011. Unpublished Scottish Natural Heritage commissioned report.

Bull C, Watt J. Site condition monitoring of lamprey in the River Teith SAC 2011. Unpublished Scottish Natural Heritage commissioned report.

Bull C, Perfect C, Watt J. 2016. Site condition monitoring of lamprey in the Endrick Water SSSI and SAC 2012. Scottish Natural Heritage Commissioned Report No. 911.

Watt J, Bull C. Site Condition Monitoring of lamprey in the River Tay SAC 2012-2013. Unpublished Scottish Natural Heritage commissioned report.

Bull C. Site condition monitoring of lamprey in the River Teith SAC 2015-2016. Unpublished Scottish Natural Heritage commissioned report.

Bull C, Shelton J, Coulson M. 2018. The collection of larval sea lamprey

(Petromyzon marinus) using drift netting, suction pumping, and airlift sampling: a trial to inform the development of methods suitable for Site Condition

Monitoring. Scottish Natural Heritage Research Report No. 1040. IAFG. 2017. UK Article 17 reporting procedure for estimating population using 1 km square resolution records data. Inter-agency Freshwater Group, UK. Records from the Atlas of Freshwater Fishes accessed through NBN Atlas website (https://registry.nbnatlas.org/public/show/dr741).

NBN Atlas website at http://www.nbnatlas.org accessed 14 June 2018.

b) Maximum

b) Maximum

5. Range

- 5.1 Surface area (km²)
- 5.2 Short-term trend Period
- 5.3 Short-term trend Direction
- 5.4 Short-term trend Magnitude
- 5.5 Short-term trend Method used
- 5.6 Long-term trend Period
- 5.7 Long-term trend Direction
- 5.8 Long-term trend Magnitude
- 5.9 Long-term trend Method used
- 5.10 Favourable reference range
- a) Area (km²)

a) Minimum

Unknown (x)

a) Minimum

- b) Operator
- c) Unknown
- d) Method

5.11 Change and reason for change in surface area of range

Improved knowledge/more accurate data Use of different method

The change is mainly due to: Use of different method

5.12 Additional information

6. Population

6.1 Year or period

1990-2018

6.2 Population size (in reporting unit)

- a) Unit
- number of map 1x1 km grid cells (grids1x1)
- b) Minimum
- c) Maximum
- d) Best single value 733

6.3 Type of estimate

Minimum

6.4 Additional population size (using population unit other than reporting unit)

- a) Unit
- b) Minimum
- c) Maximum
- d) Best single value

6.5 Type of estimate

6.6 Population size Method used

Based mainly on extrapolation from a limited amount of data

6.7 Short-term trend Period

2006-2018

6.8 Short-term trend Direction Unknown (x) 6.9 Short-term trend Magnitude a) Minimum b) Maximum c) Confidence interval 6.10 Short-term trend Method used Insufficient or no data available 6.11 Long-term trend Period 6.12 Long-term trend Direction 6.13 Long-term trend Magnitude a) Minimum b) Maximum c) Confidence interval 6.14 Long-term trend Method used 6.15 Favourable reference a) Population size population (using the unit in 6.2 or b) Operator c) Unknown d) Method 6.16 Change and reason for change Improved knowledge/more accurate data in population size Use of different method The change is mainly due to: Use of different method 6.17 Additional information 7. Habitat for the species 7.1 Sufficiency of area and quality of a) Are area and quality of occupied habitat Unknown occupied habitat sufficient (to maintain the species at FCS)? b) Is there a sufficiently large area of occupied AND unoccupied habitat of suitable quality (to maintain the species at FCS)? 7.2 Sufficiency of area and quality of Insufficient or no data available occupied habitat Method used 7.3 Short-term trend Period 2006-2018 7.4 Short-term trend Direction Unknown (x) 7.5 Short-term trend Method used Insufficient or no data available 7.6 Long-term trend Period 7.7 Long-term trend Direction 7.8 Long-term trend Method used 7.9 Additional information

8. Main pressures and threats

8.1 Characterisation of pressures/threats

Pressure Ranking

Agricultural activities generating point source pollution to M

| M |
|---------|
| М |
| М |
| M |
| Н |
| Н |
| Н |
| Н |
| Ranking |
| M |
| M |
| M |
| М |
| M |
| Н |
| Н |
| Н |
| Н |
| |

8.2 Sources of information

8.3 Additional information

9. Conservation measures

| 9.1 Status of measures | a) Are measures needed?b) Indicate the status of measures | Yes Measures identified and taken |
|----------------------------------------|--------------------------------------------------------------------------------------|------------------------------------|
| 9.2 Main purpose of the measures taken | Expand the current range of the spe | ecies (related to 'Range') |
| 9.3 Location of the measures taken | Both inside and outside Natura 2000 | 0 |

9.4 Response to the measures

Long-term results (after 2030)

9.5 List of main conservation measures

Reduce/eliminate point pollution to surface or ground waters from agricultural activities (CA10)

Reduce diffuse pollution to surface or ground waters from agricultural activities (CA11)

Reduce diffuse pollution to surface or ground waters from forestry activities (CB10)

Reduce impact of hydropower operation and infrastructure (CC04)

Reduce/eliminate point source pollution to surface or ground waters from industrial, commercial, residential and recreational areas and activities (CF04)

Reduce/eliminate diffuse pollution to surface or ground waters from industrial, commercial, residential and recreational areas and activities (CF05)

Reduce impact of multi-purpose hydrological changes (CJ02)

Adopt climate change mitigation measures (CN01)

Implement climate change adaptation measures (CN02)

9.6 Additional information

10. Future prospects

10.1 Future prospects of parameters

- a) Range
- b) Population
- c) Habitat of the species

10.2 Additional information

11. Conclusions

11.1. Range

11.2. Population

11.3. Habitat for the species

11.4. Future prospects

11.5 Overall assessment of

Conservation Status

11.6 Overall trend in Conservation

Status

11.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

11.8 Additional information

12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)

a) Unit number of map 1x1 km grid cells (grids1x1)

- b) Minimum
- c) Maximum
- d) Best single value 509

12.2 Type of estimate

12.3 Population size inside the network Method used

Minimum

Based mainly on extrapolation from a limited amount of data

12.4 Short-term trend of population size within the network Direction

Unknown (x)

12.5 Short-term trend of population size within the network Method used

Insufficient or no data available

12.6 Additional information

13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

Distribution Map




Figure 1: UK distribution map for S1095 - Sea lamprey (*Petromyzon marinus*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available species records within the current reporting period. For further details see the 2019 Article 17 UK Approach document.

Range Map




Figure 2: UK range map for S1095 - Sea lamprey (*Petromyzon marinus*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by applying a bespoke range mapping tool for Article 17 reporting (produced by JNCC) to the 10km grid square distribution map presented in Figure 1. The alpha value for this species was 25km. For further details see the 2019 Article 17 UK Approach document.

Explanatory Notes

| Species name: Petromyzon marinus (1095) | | | |
|---------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| Field label | Note | | |
| 2.2 Year or Period | Data gathered 1990-2016 have been used an exptrapolated to 2018. Only one national survey has been undertaken during this period and the results of it are still thought to be relevant to the distribution assessment. Other records gathered throughout this period are also thought to be relevant to the distribution mapping. | | |
| 2.4 Distribution map; Method used | The approach to mapping (see IAFG, 2017) is different to that used in 2013 when only actual records of the species were used to illustrate its distribution using occupied of 1 km x 1 km squares. The map produced for 2019 includes not only actual records for the species as occupied 1 km x 1 km squares but also those monads between the most upstream occupied square and the freshwater/marine interface irrespective of whether they include a record or not. | | |
| Species name: Petromyzon marinus (1095) Region code: ATL Field label Note | | | |
| 5.1 Surface area | The approach to mapping (see IAFG, 2017) is different to that used in 2013 when only actual records of the species were used to illustrate its range using occupied 1 km x 1 km squares. The map produced for 2019 includes not only actual records for the species as occupied 1 km x 1 km squares but also those monads between the most upstream occupied square and the freshwater/marine interface irrespective of whether they include a record or not. | | |
| 5.3 Short term trend; Direction | Insufficient information is available to determine the short term trend in the range of the species. To date only one national survey has been undertaken. Site Condition Monitoring surveys tend to focus on approximately the same areas. Any difference in range is likely to be the result of better information and the use of a different mapping method (see IAFG, 2017) rather than actual change. | | |
| 5.11 Change and reason for change in surface area of range | The approach to mapping (see IAFG, 2017) is different to that used in 2013 when only actual records of the species were used to illustrate its distribution using occupied of 1 km x 1 km squares. The map produced for 2019 includes not only actual records for the species as occupied 1 km x 1 km squares but also those monads between the most upstream occupied square and the freshwater/marine interface irrespective of whether they include a record or not. | | |
| 6.1 Year or Period | Data gathered 1990-2016 have been used and extrapolated to 2018. Only one national survey has been undertaken during this period and the results of it are still thought to be relevant to the distribution assessment. Other records gathered throughout this period are also thought to be relevant to the population assessment. | | |
| 6.2 Population size | The approach to estimating the population size (see IAFG, 2017) is different to that used in 2013 when only actual records of occupied 1 km x 1 km squares were used. The estimate produced for 2019 includes not only actual records of 1 km x 1 km squares occupied by the species but also those monads between the most upstream and and the freshwater/marine interface irrespective of whether they include a record or not. The best single value for occupied wholly Scotland 1 km x 1 km squares is 733. In addition there are 39 occupied Scotland/England transboundary 1 km x 1 km squares. | | |
| 6.8 Short term trend; Direction | Insufficient information is available to determine the short term trend in the population of the species. Any attempt to determine what the trend might be would be complicated by the transient nature of the species' habitat and interannual variation in the number of individuals. | | |

| 6.16 Change and reason for change in population size | The change in population size reflects the new method (see IAFG, 2017) of mapping the species, i.e. actual records for the species as occupied 1 km x 1 km squares and the 1 km x 1 km squares between the most upstream records and the freshwater/marine interface, irrespective of whether they include a record or not, have been included. Assessments of the true population size would be unlikely to be meaningful due to the frequency of surveys, the transient nature of the species' habitat, and interannual variation in the number of individuals. |
|----------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 7.1 Sufficiency of area and quality of occupied habitat | Surveys focus on the larvae and historically the same technique has been applied to all three lamprey species. The number of Petromyzon marinusrecorded during surveys is low and frequently two orders of magnitude less than the number of Lampetra. Low numbers of P. marinusmay be being caught for a variety of reasons including: they are naturally relatively scarce; they have different habitat preferences and their favoured habitat is not being adequately surveyed; and access to their habitat is hampered by engineered structures, e.g. weirs, dams. The sufficiency of area and quality of occupied habitat remains unknown. |
| 8.1 Characterisation of pressures/ threats | Pressures arising from inadequate land management and pollution control and river engineering are likely to persist in the short to medium term. Regulation aimed at dealing with these issues exits and is evolving but there may be a lag in the biological response to improvements. The exact nature of the direct and indirect effects of climate change remain unknown. Pressures on the marine environment may affect sea lamprey but insufficient is known about the adult phase. |
| 9.1 Status of measures | The measures given in the 2013 report were some of those needed to ensure that sea lamprey habitat was maintained or restored. However, non was undertaken specifically for sea lamprey. The situation is the same for the current reporting period. The presumption is that the presence of good habitat is likely to result in healthy populations. Research undertaken in the intervening period suggests that not enough is known about the species to be able to specify its exact habitat requirements and so what, if anything, might be adversely affecting them. Meaningful habitat and population assessments are needed to identify species specific conservation measures. In order for these to be undertaken better information about the species including its life history is needed. |
| 10.1 Future prospects of parameters | Reliable information about the range, population, and habitat of the species has yet to be established. Until it is it will not be possible to provide a meaningful prediction of the prospects for these parameters. Research into new lamprey survey and identification techniques (notably the use of eDNA) has been undertaken in recent years but further work is needed. |
| 12.1 Population size inside the pSCIs, SCIs and SACs network | The best single value for occupied wholly Scotland 1 km x 1 km squares is 509. In addition there are 39 occupied Scotland/England transboundary 1 km x 1 km squares. |
| 12.4 Short term trend of the population size within the network; Direction | Insufficient information is available to determine the short term trend in the population within the network. Any attempt to determine what the trend might be would be complicated by the transient nature of the species' habitat and interannual variation in the number of individuals. |