

**European Community Directive
on the Conservation of Natural Habitats
and of Wild Fauna and Flora
(92/43/EEC)**

**Fourth Report by the United Kingdom
under Article 17**

on the implementation of the Directive
from January 2013 to December 2018

Supporting documentation for the
conservation status assessment for the species:

S1309 - Common pipistrelle (*Pipistrellus pipistrellus*)

SCOTLAND

IMPORTANT NOTE - PLEASE READ

- The information in this document is a country-level contribution to the UK Report on the conservation status of this species, submitted to the European Commission as part of the 2019 UK Reporting under Article 17 of the EU Habitats Directive.
- The 2019 Article 17 UK Approach document provides details on how this supporting information was used to produce the UK Report.
- The UK Report on the conservation status of this species is provided in a separate document.
- The reporting fields and options used are aligned to those set out in the European Commission guidance.
- Explanatory notes (where provided) by the country are included at the end. These provide an audit trail of relevant supporting information.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; (iii) the field was not relevant to this species (section 12 Natura 2000 coverage for Annex II species) and/or (iv) the field was only relevant at UK-level (sections 9 Future prospects and 10 Conclusions).
- For technical reasons, the country-level future trends for Range, Population and Habitat for the species are only available in a separate spreadsheet that contains all the country-level supporting information.
- The country-level reporting information for all habitats and species is also available in spreadsheet format.

Visit the JNCC website, <https://jncc.gov.uk/article17>, for further information on UK Article 17 reporting.

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

NATIONAL LEVEL

1. General information

| | |
|---|--------------------------------|
| 1.1 Member State | UK (Scotland information only) |
| 1.2 Species code | 1309 |
| 1.3 Species scientific name | Pipistrellus pipistrellus |
| 1.4 Alternative species scientific name | |
| 1.5 Common name (in national language) | Common pipistrelle |

2. Maps

| | |
|----------------------------------|---|
| 2.1 Sensitive species | No |
| 2.2 Year or period | 1995-2016 |
| 2.3 Distribution map | Yes |
| 2.4 Distribution map Method used | Based mainly on extrapolation from a limited amount of data |
| 2.5 Additional maps | No |

3. Information related to Annex V Species (Art. 14)

| | | | | | | | | | | | | | | | | | |
|---|--|---|----|---|----|---|----|--|----|--|----|---|----|---|----|-------------------|----|
| 3.1 Is the species taken in the wild/exploited? | No | | | | | | | | | | | | | | | | |
| 3.2 Which of the measures in Art. 14 have been taken? | <table> <tr> <td>a) regulations regarding access to property</td><td>No</td></tr> <tr> <td>b) temporary or local prohibition of the taking of specimens in the wild and exploitation</td><td>No</td></tr> <tr> <td>c) regulation of the periods and/or methods of taking specimens</td><td>No</td></tr> <tr> <td>d) application of hunting and fishing rules which take account of the conservation of such populations</td><td>No</td></tr> <tr> <td>e) establishment of a system of licences for taking specimens or of quotas</td><td>No</td></tr> <tr> <td>f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens</td><td>No</td></tr> <tr> <td>g) breeding in captivity of animal species as well as artificial propagation of plant species</td><td>No</td></tr> <tr> <td>h) other measures</td><td>No</td></tr> </table> | a) regulations regarding access to property | No | b) temporary or local prohibition of the taking of specimens in the wild and exploitation | No | c) regulation of the periods and/or methods of taking specimens | No | d) application of hunting and fishing rules which take account of the conservation of such populations | No | e) establishment of a system of licences for taking specimens or of quotas | No | f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens | No | g) breeding in captivity of animal species as well as artificial propagation of plant species | No | h) other measures | No |
| a) regulations regarding access to property | No | | | | | | | | | | | | | | | | |
| b) temporary or local prohibition of the taking of specimens in the wild and exploitation | No | | | | | | | | | | | | | | | | |
| c) regulation of the periods and/or methods of taking specimens | No | | | | | | | | | | | | | | | | |
| d) application of hunting and fishing rules which take account of the conservation of such populations | No | | | | | | | | | | | | | | | | |
| e) establishment of a system of licences for taking specimens or of quotas | No | | | | | | | | | | | | | | | | |
| f) regulation of the purchase, sale, offering for sale, keeping for sale or transport for sale of specimens | No | | | | | | | | | | | | | | | | |
| g) breeding in captivity of animal species as well as artificial propagation of plant species | No | | | | | | | | | | | | | | | | |
| h) other measures | No | | | | | | | | | | | | | | | | |

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

3.3 Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

| b) Statistics/ quantity taken | Provide statistics/quantity per hunting season or per year (where season is not used) over the reporting period | | | | | |
|----------------------------------|---|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Season/ year 1 | Season/ year 2 | Season/ year 3 | Season/ year 4 | Season/ year 5 | Season/ year 6 |
| Min. (raw, ie. not rounded) | | | | | | |
| Max. (raw, ie. not rounded) | | | | | | |
| Unknown | No | No | No | No | No | No |

3.4. Hunting bag or quantity taken in the wild Method used

3.5. Additional information

BIOGEOGRAPHICAL LEVEL

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs

Atlantic (ATL)

4.2 Sources of information

Barratt, E.M., R. Deaville, T.M. Burland, M.W. Bruford, G. Jones, P.A. Racey and R.K. Wayne (1997). DNA answers the call of pipistrelle bat species. *Nature*, 387 (6629), 138-139

Bat Conservation Trust (2018). The State of the UK's Bats 2017. Bat Conservation Trust, London. Available at (http://www.bats.org.uk/pages/results_and_reports.html)

Boye, P and Dietz, M. (2005). Research Report No 661: Development of good practice guidelines for woodland management for bats. English Nature, Peterborough.

Davidson-Watts, I. and Jones, G. (2006). Differences in foraging behaviour between *Pipistrellus pipistrellus* (Schreber, 1774) and *Pipistrellus pygmaeus* (Leach, 1825). *Journal of Zoology*, 268 (1), 55-62

Fensome, A. G. and Mathews, F. (2016). Roads and bats: a meta-analysis and review of evidence on vehicle collisions and barrier effects. *Mammal Review*, 46 (4), 311-323

Fuentes-Montemayor, E., Goulson, D., Cavin, L., Wallace, J.M., and Park, K.J. (2013). Fragmented woodlands in agricultural landscapes: The influence of woodland character and landscape context on bats and their insect prey. *Agriculture, Ecosystems and Environment*, 172, 6-15

Glendell, M. and Vaughan, N. (2002). Foraging activity of bats in historic landscape parks in relation to habitat composition and park management. *Animal Conservation*, 5 (4), 309-316

Jones, G and Racey, P.A. (2008). Common pipistrelle *Pipistrellus pipistrellus*, Soprano pipistrelle *Pipistrellus pygmaeus*. Pages 343-351 In Harris, S and Yalden, D.W. *Mammals of the British Isles: Handbook*, 4th edition. The Mammal Society,

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

Southampton. 799pp.

Lintott, P.R., Bunnefeld, N. and Park, K.J. (2015). Opportunities for improving the foraging potential of urban waterways for bats. *Biological Conservation*, 191, 224-233.

Lintott, P.R., Barlow, K., Bunnefeld, N., Briggs, P., Gajas Roig, C., and Park, K.J. (2016). Differential responses of cryptic bat species to the urban landscape. *Ecology and Evolution*, 6 (7), 2044-2052

Mathews, F., Richardson, S.M., and Hosken, D.J. (2016). Understanding the risks to bat populations posed by wind turbines - Phase 2 - WC0753, Defra.

Mathews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C., McDonald, R.A., Shore, R.F (2018). A review of the population and conservation status of British Mammals. A report by the Mammal Society under contract to Natural England, Natural Resources Wales and Scottish Natural Heritage.

Mitchell-Jones, T.J. (2010). Bats in houses - the conservation challenge. Pp 3965-378 in *Species Management : challenges and solutions for the 21st century*.

Baxter, J.M. and Galbraith, C.A. TSO Scotland, Edinburgh

Newson, S.E., Evans, H.E., Gillings, S., Jarrett, D. & Wilson, M.W. 2017. A survey of high risk bat species across southern Scotland. Scottish Natural Heritage Commissioned Report No. 1008.

Nicholls, B. and Racey, P (2006a). Habitat selection as a mechanism of resource partitioning in two cryptic bat species *Pipistrellus pipistrellus* and *pipistrellus pygmaeus*. *Ecography*, 29, 697-708

Nicholls, B. and Racey, P (2006b). Contrasting home-range size and spatial partitioning in cryptic and sympatric bats. *Behavioural Ecology and Sociobiology*, 61, 131-142

Waring, S.D., Essah, E., Gunnell, K, and Bonser, R (2013). Double jeopardy: the potential for problems when bats interact with breathable roofing membranes in the United Kingdom. *Architecture and Environment*, 1 1-3

Warren, R, D., Waters, D, A., Altringham, J.D., and Bullock, D.J. (2000). The distribution of Daubenton's bats (*Myotis daubentonii*) and pipistrelle bats (*Pipistrellus pipistrellus*) (*Vespertilionidae*) in relation to small-scale variation in riverine habitat. *Biological Conservation*, 92 (1), 85-91

5. Range

5.1 Surface area (km²)

5.2 Short-term trend Period

5.3 Short-term trend Direction

Stable (0)

5.4 Short-term trend Magnitude

a) Minimum

b) Maximum

5.5 Short-term trend Method used

5.6 Long-term trend Period

5.7 Long-term trend Direction

5.8 Long-term trend Magnitude

a) Minimum

b) Maximum

5.9 Long-term trend Method used

5.10 Favourable reference range

a) Area (km²)

b) Operator

c) Unknown

d) Method

5.11 Change and reason for change in surface area of range

No change

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

The change is mainly due to:

5.12 Additional information

6. Population

| | | | |
|--|---|--|--|
| 6.1 Year or period | 2016-2017 | | |
| 6.2 Population size (in reporting unit) | a) Unit | number of map 1x1 km grid cells (grids1x1) | |
| | b) Minimum | | |
| | c) Maximum | | |
| | d) Best single value | | |
| 6.3 Type of estimate | Minimum | | |
| 6.4 Additional population size (using population unit other than reporting unit) | a) Unit | number of individuals (i) | |
| | b) Minimum | 285000 | |
| | c) Maximum | 2160000 | |
| | d) Best single value | | |
| 6.5 Type of estimate | 95% confidence interval | | |
| 6.6 Population size Method used | Based mainly on extrapolation from a limited amount of data | | |
| 6.7 Short-term trend Period | 2007-2018 | | |
| 6.8 Short-term trend Direction | Stable (0) | | |
| 6.9 Short-term trend Magnitude | a) Minimum | | |
| | b) Maximum | | |
| | c) Confidence interval | | |
| 6.10 Short-term trend Method used | Based mainly on extrapolation from a limited amount of data | | |
| 6.11 Long-term trend Period | | | |
| 6.12 Long-term trend Direction | | | |
| 6.13 Long-term trend Magnitude | a) Minimum | | |
| | b) Maximum | | |
| | c) Confidence interval | | |
| 6.14 Long-term trend Method used | | | |
| 6.15 Favourable reference population (using the unit in 6.2 or 6.4) | a) Population size | | |
| | b) Operator | | |
| | c) Unknown | | |
| | d) Method | | |
| 6.16 Change and reason for change in population size | Improved knowledge/more accurate data | | |
| | Use of different method | | |
| | The change is mainly due to: | Use of different method | |
| 6.17 Additional information | | | |

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

7. Habitat for the species

| | | |
|---|--|-----|
| 7.1 Sufficiency of area and quality of occupied habitat | a) Are area and quality of occupied habitat sufficient (to maintain the species at FCS)? b) Is there a sufficiently large area of occupied AND unoccupied habitat of suitable quality (to maintain the species at FCS)? | Yes |
| 7.2 Sufficiency of area and quality of occupied habitat Method used | Based mainly on extrapolation from a limited amount of data | |
| 7.3 Short-term trend Period | 2007-2018 | |
| 7.4 Short-term trend Direction | Stable (0) | |
| 7.5 Short-term trend Method used | Based mainly on extrapolation from a limited amount of data | |
| 7.6 Long-term trend Period | | |
| 7.7 Long-term trend Direction | | |
| 7.8 Long-term trend Method used | | |
| 7.9 Additional information | | |

8. Main pressures and threats

8.1 Characterisation of pressures/threats

| Pressure | Ranking |
|---|---------|
| Use of plant protection chemicals in agriculture (A21) | M |
| Conversion to other types of forests including monocultures (B02) | M |
| Logging without replanting or natural regrowth (B05) | M |
| Wind, wave and tidal power, including infrastructure (D01) | M |
| Roads, paths, railroads and related infrastructure (e.g. bridges, viaducts, tunnels) (E01) | H |
| Construction or modification (e.g. of housing and settlements) in existing urban or recreational areas (F02) | H |
| Residential or recreational activities and structures generating noise, light, heat or other forms of pollution (F24) | H |
| Industrial or commercial activities and structures generating noise, light, heat or other forms of pollution (F25) | H |
| Removal of small landscape features for agricultural land parcel consolidation (hedges, stone walls, rushes, open ditches, springs, solitary trees, etc.) (A05) | H |
| Threat | Ranking |
| Use of plant protection chemicals in agriculture (A21) | M |
| Conversion to other types of forests including monocultures (B02) | M |
| Logging without replanting or natural regrowth (B05) | M |
| Wind, wave and tidal power, including infrastructure (D01) | M |

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

Roads, paths, railroads and related infrastructure (e.g. bridges, viaducts, tunnels) (E01) H

Construction or modification (e.g. of housing and settlements) in existing urban or recreational areas (F02) H

Residential or recreational activities and structures generating noise, light, heat or other forms of pollution (F24) H

Industrial or commercial activities and structures generating noise, light, heat or other forms of pollution (F25) H

Removal of small landscape features for agricultural land parcel consolidation (hedges, stone walls, rushes, open ditches, springs, solitary trees, etc.) (A05) H

8.2 Sources of information

8.3 Additional information

9. Conservation measures

9.1 Status of measures

a) Are measures needed? Yes

b) Indicate the status of measures Measures identified and taken

9.2 Main purpose of the measures taken

Maintain the current range, population and/or habitat for the species

9.3 Location of the measures taken

Both inside and outside Natura 2000

9.4 Response to the measures

Medium-term results (within the next two reporting periods, 2019-2030)

9.5 List of main conservation measures

Prevent conversion of natural and semi-natural habitats, and habitats of species into agricultural land (CA01)

Other measures related to agricultural practices (CA16)

Prevent conversion of (semi-) natural habitats into forests and of (semi-)natural forests into intensive forest plantation (CB01)

Adapt/manage reforestation and forest regeneration (CB04)

Adapt/manage renewable energy installation, facilities and operation (CC03)

Reduce impact of transport operation and infrastructure (CE01)

Reduce/eliminate noise, light, heat or other forms pollution from industrial, commercial, residential and recreational areas and activities (CF09)

Reduce/eliminate diffuse pollution to surface or ground waters from industrial, commercial, residential and recreational areas and activities (CF05)

Restore small landscape features on agricultural land (CA02)

9.6 Additional information

10. Future prospects

10.1 Future prospects of parameters

- a) Range
- b) Population
- c) Habitat of the species

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

10.2 Additional information

11. Conclusions

11.1. Range

11.2. Population

11.3. Habitat for the species

11.4. Future prospects

11.5 Overall assessment of Conservation Status

11.6 Overall trend in Conservation Status

11.7 Change and reasons for change in conservation status and conservation status trend

a) Overall assessment of conservation status

No change

The change is mainly due to:

b) Overall trend in conservation status

No change

The change is mainly due to:

11.8 Additional information

12. Natura 2000 (pSCIs, SCIs and SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network (on the biogeographical/marine level including all sites where the species is present)

- a) Unit
- b) Minimum
- c) Maximum
- d) Best single value

12.2 Type of estimate

12.3 Population size inside the network Method used

12.4 Short-term trend of population size within the network Direction

12.5 Short-term trend of population size within the network Method used

12.6 Additional information

13. Complementary information

13.1 Justification of % thresholds for trends

13.2 Trans-boundary assessment

13.3 Other relevant Information

Report on the main results of the surveillance under Article 11 for Annex II, IV and V species (Annex B)

Distribution Map




Figure 1: UK distribution map for S1309 - Common pipistrelle (*Pipistrellus pipistrellus*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The 10km grid square distribution map is based on available species records within the current reporting period. For further details see the 2019 Article 17 UK Approach document.

Range Map




Figure 2: UK range map for S1309 - Common pipistrelle (*Pipistrellus pipistrellus*). Coastline boundary derived from the Oil and Gas Authority's OGA and Lloyd's Register SNS Regional Geological Maps (Open Source). Open Government Licence v3 (OGL). Contains data © 2017 Oil and Gas Authority.

The range map has been produced by The Mammal Society applying a range mapping tool as outlined in Matthews et al. (2018), to the 10km grid square distribution map presented in Figure 1. The alpha value for this species was 20km. For further details see the 2019 Article 17 UK Approach document.

Explanatory Notes

Species name: Pipistrellus pipistrellus (1309) Region code: ATL

| Field label | Note |
|---|---|
| 5.3 Short term trend; Direction | Range is based on presence data collected between 1995-2016. Areas that contain very isolated records may not have been included in the area of distribution. The range has been taken from Mathews et al 2018, whereby an alpha hull value of 20km was drawn around the presence records, which represented the best balance between the inclusion of unoccupied sites (i.e. where records are sparse but close enough for inclusion) and the exclusion of occupied areas due to gaps in the data (i.e. where records exist but are too isolated for inclusion). An additional 10km buffer was added to the final hull polygon to provide smoothing to the hull and to ensure that the hull covered the areas recorded rather than intersecting them. This differs from the approach taken in 2013 and 2007 whereby a 45km alpha hull value was used for all species with a starting range unit of individual 10km squares. The new method has led to much finer detail maps being produced underpinned by data gathered at a much finer resolution. The stable trend is based on expert opinion and the Mathews et al. range map. |
| 5.10 Favourable reference range | Range is based on presence data collected between 1995-2016. Areas that contain very isolated records may not have been included in the area of distribution. The range has been taken from Mathews et al 2018, whereby an alpha hull value of 20km was drawn around the presence records, which represented the best balance between the inclusion of unoccupied sites (i.e. where records are sparse but close enough for inclusion) and the exclusion of occupied areas due to gaps in the data (i.e. where records exist but are too isolated for inclusion). An additional 10km buffer was added to the final hull polygon to provide smoothing to the hull and to ensure that the hull covered the areas recorded rather than intersecting them. This differs from the approach taken in 2013 and 2007 whereby a 45km alpha hull value was used for all species with a starting range unit of individual 10km squares. The new method has led to much finer detail maps being produced underpinned by data gathered at a much finer resolution, leading to the production of a more accurate FRR. Added to which acoustic detectors have changed considerably over the years in both accuracy and sensitivity, which also adds to the production of this value. |
| 6.4 Additional population size | Mathews et al, (2018) gives estimates of 285,000 individuals (lower plausible limit) to 2,160,000 (upper plausible limit). The estimates excluded colonies that contained less than 30 bats in order to ensure that counts did not include individuals in formation roosts that were then counted again at maternity sites. This may have led to some over-estimation of population size: when all roosts were included the bat population density estimate fell by approximately a third. However, most data were derived from NBMP data and here all roosts were included regardless of size since they were part of a longitudinal monitoring programme. Given that the estimated roost size is close to expert opinion and published data, it is likely to be a reasonable basis for the calculations. |
| 6.8 Short term trend; Direction | The National Bat Monitoring Programme (NBMP) population trends record no significant change in the Scottish population since the last Article 17 reporting round. There would seem to have been a significant decline in the short-term population trend for this species from roost counts. However, it is likely that for this species, frequent roost switching results in a negative bias in Roost Count data. The Roost Count trend is therefore not considered a reliable measure of population change for this species. |
| 6.16 Change and reason for change in population size | Also improved knowledge. Acoustic detectors used to record bat activity in the field have changed considerably over time and have become much more sensitive. |

| | |
|---|---|
| 7.1 Sufficiency of area and quality of occupied habitat | <p><i>P. pipistrellus</i> is an extremely widespread species and is found in almost any habitat type ranging from grasslands to urban and suburban environments. However, the species requires a complex mosaic of habitats to support foraging, roosting and commuting behaviour. Boye & Dietz (2005) and Jones & Racey (2008) provides a good overview of this species habitat requirements. Although, most maternity colonies are in buildings, forests of any type are used as roosting and foraging areas. Foraging areas are mainly along woodland edge and riparian woodland (Davidson-Watts & Jones 2006; Nicholls & Racey 2006a, 2006b), hedges, foot paths and forest roads, water banks and at street lights. <i>P. pipistrellus</i> frequently forage over pasture and foraging activity is higher where grazing livestock are present (Fuentes-Montemayor et al 2013). Linear features in a landscape are important elements for orientation either during foraging or in commuting flights. Foraging activity is generally within 2km of the roost. The size of an individual home range is dependent on the abundance of prey insects and may have a total size of more than 50 hectares. The species mainly roosts in settlements and is even present in city centres. Recent evidence shows that there is a strong negative response of <i>P. pipistrellus</i> to urbanisation at a relatively local scale (1km; Lintott et al 2016). However, the reverse association has also been reported (Warren et al. 2000, Glendell & Vaughan 2002, Lintott et al. 2015). In summer the roost sites are predominantly in crevices in buildings, especially between tiles and the underlying roofing felt or behind boards on the gable. Furthermore, individuals and maternity colonies use tree holes, wood crevices and bird or bat boxes as roosts. The species disperses to temporary sites and mating roosts during the autumn post weaning period. As demonstrated by the species' extensive range and overall abundance, there is sufficient habitat in Scotland to support a viable population of the species.</p> |
| 7.4 Short term trend; Direction | <p>This stable habitat trend is expert opinion, with Mathews et al suggesting that the habitat future prospects for this species is stable.</p> |
| 8.1 Characterisation of pressures/ threats | <p>Pressures can generally be divided into those that affect roosts and those that affect commuting and foraging (including prey availability). <i>P. pipistrellus</i> forage across a mosaic of habitat types, though they are frequently found foraging over pasture, especially at sites with grazing livestock (Fuentes-Montemayor et al, 2013). Agricultural and forestry practices that remove or simplify these habitats or affect the biomass of insect prey could negatively affect populations. This is one of the primary species killed at wind turbine sites and in road collisions. It is unclear whether the scale of casualties is sufficient to impact on local populations (Mathews et al, 2016 and Fensome & Mathews, 2016). Although, roosts are strictly protected through legislation a variable number of licences are issued every year permitting exclusion, destruction and damage. Changes to building regulations and efforts to make buildings more energy-efficient have tended to reduce their accessibility and thermal suitability for bats.</p> |
| 9.5 List of main conservation measures | <p>Legal and administrative measures continue to be required to ensure that the protection provided by the legislation is effective. If roosts are to be destroyed, damaged or lost due to development, adequate mitigation/compensation methods must be put in place to maintain the favourable conservation status of the species. Road design construction and operation need to take into account the likely impact on bats, for example, in relation to the provision of safe crossing structures and the loss and severance of bat habitat and lighting. Guidance is being developed and will shortly be available from the agencies to help planners, developers and ecological consultants to consider the potential effects of onshore wind energy developments on bats. Guidance is available for land managers on how to manage their land holdings for bats.</p> |

10.1 Future prospects of parameters

The range for *P. pipistrellus* is likely to remain stable as the species continues to be widespread and appears to be covering roughly the same range as in the previous reporting round (2007-2012), even though different methods were used to perform this calculation. The population in Scotland is considered to be stable (NBMP trend data). It is considered that the habitat for *P. pipistrellus* will remain stable: it is taken as being related to the range as the species is widespread occupying a wide variety of habitats and habitat mosaics.
